Michigan General Procedures PATIENT RESTRAINT

Date: May 31, 2012 Page 1 of 2

Patient Restraint

Purpose: To ensure appropriate restraint of patients and to assure patient, others and EMS safety.

Pre-Medical Control

MFR/EMT/SPECIALIST/PARAMEDIC

Indications:

1. When an ill or injured person who is behaving in such a manner as to interfere with their examination, care and treatment to the extent they endanger their life or the safety of others.

Physical Restraint Procedure

- 1. Ensure that enough personnel are available to properly control the patient and establish the restraints.
- 2. Explain the purpose of the restraints.
- 3. Physically control the patient and apply restraints.
- 4. Complete Primary and Secondary Assessments.
 - A. Restrained extremities should be evaluated for pulse quality, capillary refill time, color, sensory and motor function continuously
 - a. Restraints must be adjusted if any of these functions are compromised.
 - b.Restraints must not interfere with medical treatment.
- 5. Attempt to identify common physical causes for patient's abnormal behavior.
 - Hypoxia
 - Hypoglycemia
 - Head Trauma
 - ETOH/ Substances use/ abuse
- 6. Patient should be secured to a backboard or stretcher only. Patients must never be secured directly to a vehicle or immovable object.
- 7. Transport patient.
- 8. Contact medical control.
- 9. Inform hospital that restraints are in place and assistance will be necessary to continue restraint of the patient.

Post-Medical Control

PARAMEDIC

Chemical Restraint Procedure

- 1. If Chemical restraint is considered, contact medical control for appropriate guidance; also refer to **Patient Sedation Procedure.**
- 2. Chemical restraint may only be performed under direct medical control order.

Special Considerations

1. Physical restraints should be of a soft nature (e.g. leather cuffs, cravats, sheets, etc.) applied to the wrists and ankles. A restraint may also be needed across the chest and/or pelvis.



Michigan

General Procedures PATIENT RESTRAINT

Date: May 31, 2012 Page 2 of 2

- 2. Stay with a restrained patient at all times, be observant for possible vomiting and be prepared to turn the patient and suction if necessary.
- 3. Documentation should include:
 - A. A description of the circumstance / behavior which precipitated the use of restraints.
 - B. Time of application of the restraints.
 - C. Type of restraint used.
 - D. The positions in which the patient was restrained.
- 4. When restraint devices are applied by law enforcement officers:
 - A. An officer must be present with the patient at all times at the scene, as well as in the ambulance during transport.
 - B. The restraint and position must not be so restrictive that the patient is in a position that compromise patient care.
- 5. EMS Personnel may NOT use:
 - A. Hard plastic ties or any restraint devices that require a key to remove.
 - B. Backboards to "sandwich" the patient.
 - C. Restraints which secures the patient's hands and feet behind the back.
 - D. Restraints that "hog tie" the patient.
 - E. Any device that restricts normal breathing.

