

## ***Pediatric Altered Mental Status***

The purpose of this protocol is to provide for the assessment and treatment of pediatric patients with altered mental status of unknown etiology such as alcohol, trauma, poisonings, seizures, behavioral problems, stroke, environmental causes, infection, etc.

- For pediatrics less than < 24 hours old – refer to **Newborn/Neonatal Assessment and Resuscitation-Treatment Protocol**
- For critically ill patients refer to **Pediatric Crashing Patient/Impending Arrest-Treatment Protocol**
- 1. Follow **General Pre-hospital Care-Treatment Protocol**.
- 2. Pediatric patients ( $\leq 14$  years) utilize MI MEDIC cards for appropriate medication dosage. When unavailable utilize pediatric dosing listed within protocol
- 3. Restrain patient, if necessary, refer to **Patient Restraint-Procedure Protocol**.
- 4. Ensure adequate oxygenation, ventilation, and work of breathing
  - Ⓜ A. Monitor SpO<sub>2</sub>
  - Ⓢ B. Consider use of capnography
- Ⓜ 5. Check blood glucose (may be MFR skill, see **Blood Glucose Testing-Procedure Protocol**)
- 6. Check temperature if febrile go to **Pediatric Fever-Treatment Protocol**
- Ⓢ 7. Start IV/IO if needed per **Vascular Access & IV Therapy-Procedure Protocol**
- 8. Altered and able to swallow – administer **oral glucose** if:
  - A. 2 months old or younger and glucose is <40 mg/dL
  - B. 3 months old or older and glucose is <60 mg/dL .
- Ⓢ 9. Not alert – administer **dextrose** according to MI-MEDICS CARDS or table below
  - A. 2 months old or younger and glucose is <40 mg/dL
  - B. 3 months old or older and glucose is <60 mg/dL

Color	Age	Weight	Dose	Concentration	Volume		Concentration	Volume
Grey	0-2 months	3-5 kg (6-11 lbs.)	2.5g	Dextrose 12.5%	20 mL	<b>OR</b>	Dextrose 10%	25 mL
Pink	3-6 months	6-7 kg (13-16 lbs.)	3.25g	Dextrose 25%	13 mL	<b>OR</b>	Dextrose 10%	33 mL
Red	7-10 months	8-9 kg (17-20 lbs.)	4.25g	Dextrose 25%	17 mL	<b>OR</b>	Dextrose 10%	43 mL
Purple	11-18 months	10-11 kg (21-25 lbs.)	5g	Dextrose 25%	20 mL	<b>OR</b>	Dextrose 10%	50 mL
Yellow	19-35 months	12-14 kg (26-31 lbs.)	6.25g	Dextrose 25%	25 mL	<b>OR</b>	Dextrose 10%	63 mL
White	3-4 years	15-18 kg (32-40 lbs.)	8g	Dextrose 25%	32 mL	<b>OR</b>	Dextrose 10%	80 mL
Blue	5-6 years	19-23 kg (41-50 lbs.)	10g	Dextrose 25%	40 mL	<b>OR</b>	Dextrose 10%	100 mL
Orange	7-9 years	24-29 kg (52-64 lbs.)	12.5g	Dextrose 50%	25 mL	<b>OR</b>	Dextrose 10%	125 mL
Green	10-14 Years	30-36 kg (65-79 lbs.)	15g	Dextrose 50%	40 mL	<b>OR</b>	Dextrose 10%	150 mL

Initial Date: 11/2012

Revised Date: 05/24/2023

Section: 4-4

- ① 10. Per MCA selection, if unable to start IV, administer **glucagon** IM/IN (if available per MCA selection) according to MI-MEDIC cards, (may be EMT skill per MCA selection). If MI MEDIC cards are unavailable following dosing as below.

**Glucagon administration per MCA Selection**

Not included

	<u><b>Glucagon IM</b></u>	<u><b>Glucagon IN</b></u>
	A. Patients less than 5 years of age administer <b>glucagon</b> 0.5 mg IM	A. Patients less than 5 years of age, administer <b>glucagon</b> 0.5 mg IN
	B. Patients aged 5 or greater, administer <b>glucagon</b> 1 mg IM	B. Patients aged 5 or greater, administer <b>glucagon</b> 1 mg IN
EMT	<input type="checkbox"/>	<input type="checkbox"/>
Specialist	<input type="checkbox"/>	<input type="checkbox"/>
Paramedic	<input type="checkbox"/>	<input type="checkbox"/>

11. If patient respiratory depression persists and/or patient has not regained consciousness despite adequate oxygenation and ventilatory support administer **naloxone** per **Opioid Overdose Treatment and Prevention-Treatment Protocol**



12. Contact Medical Control for repeat **dextrose**.



13. Contact Medical Control for repeat **naloxone**.

**NOTE:**

- Instructions for diluting **dextrose**
  - To obtain **dextrose 10%**, discard 40 ml out of one amp of D50, then draw up 40 ml of **NS** into the D50 ampule
  - To obtain **dextrose 12.5%**, discard 37.5 ml out of one amp of D50, then draw 37.5 ml of **NS** into the D50 ampule.
  - To obtain **dextrose 25%**, discard 25 ml out of one amp of D50, then draw 25 ml of **NS** into the D50 ampule.

b. May utilize **dextrose 10%** for all ages 5 ml/kg (0.5 gm/kg) up to 250 ml, according to **Dextrose-Medication Protocol**.
- To avoid extravasation, a patent IV must be available for IV administration of **dextrose**. **Dextrose** should always be pushed slowly (e.g., over 1-2 minutes).

Medication Protocols

Dextrose

Glucagon

Naloxone