

**HEMS**  
**MEDICATION SECTION (INTERFACILITY TRANSFER)**  
**GLYCOPROTEIN IIb/IIIa RECEPTOR ANTAGONIST FOR ACUTE CORONARY SYNDROMES**

Date: March 29, 2019

Section 9.54

---

***Glycoprotein IIb/IIIa Receptor Antagonist for Acute Coronary Syndromes***

**Indications:**

In combination with Heparin for the treatment of acute coronary syndromes, including unstable angina and non Q-wave AMI and for those patients who are to be managed medically and those undergoing PTCA or artherectomy.

**Contraindications:**

1. Known hypersensitivity.
2. Active internal bleeding or a history of significant bleeding within the previous 30 days.
3. History of intracranial hemorrhage, intracranial neoplasm, arteriovenous malformation or aneurysm.
4. History of thrombocytopenia following prior exposure to Aggrastat, Integrelin or ReoPro.
5. History of CVA within 30 days or any history of hemorrhagic stroke.
6. Major surgical procedure or severe physical trauma within previous month.
7. History, symptoms or findings suggestive of aortic dissection.
8. Severe HTN (systolic BP > 180 mmHg and/or diastolic BP > 110 mmHg).
9. Concomitant use of another parenteral IIb/IIIa inhibitor.

**Procedure:**

1. Maintain appropriate delivery rate by IV infusion pump,
2. Monitor V/S every 5-10 min.

**Adverse Effects:**

Bleeding, edema/swelling, hypotension, bradycardia, pain, dizziness, sweating, nausea.

If patient develops severe hypotension, active bleeding or hypersensitivity discontinue and notifying sending facility. If sending facility is not available contact medical control.

**\*\*Not in Medication Box**