

HEMS
MEDICATION SECTION
2-PAM CHLORIDE/DUODOTE
(Protopam Chloride/
Pralidoxime)

Initial Date: 10/25/2017
Revised Date:

Section 9-55

2-Pam Chloride/DuoDote

Protocols:

1. Nerve Agent Organophosphate exposure







Indications:

1. Exposure to organophosphate or nerve agents
2. Given in conjunction with atropine in DuoDote or Mark-1 kit

Contraindications:

1. None

Dosing:

1. Self-Rescue – 1 DuoDote (Mark-1) Injector
2. Mild Reaction
 - a. Adults (8 years and over) – 1 DuoDote (Mark-1) Injector
 -   b. Pediatrics – Contact Medical Control
3. Moderate Reaction
 - a. Adults (8 years and over) – 2 DuoDote (Mark-1) Injectors
 -   b. Pediatrics – Contact Medical Control
4. Severe Reaction
 - a. Adults (8 years and over) – 3 DuoDote (Mark-1) Injector
 -   b. Pediatrics – 1 DuoDote (Mark-1) Injector, Contact Medical Control as needed

Expected Effects:

1. Decrease in symptoms

Side Effects:

1. Blurred vision
2. Headache
3. Dizziness
4. Nausea

****Not in Medication Box**