

Initial Date: 07/19/2023
Revised Date: 02/10/2026

Section: 9-35R

Morphine

Pharmacological Category: Analgesic, Opioid

Routes: IV/IO/IM

Indications:

1. Pain

Precautions:

1. Pregnant or suspected pregnant patients
 - a. Optimize non-pharmacologic interventions as directed in the **Pain Management Protocol**.
 - b. For mild to moderate pain, use acetaminophen as directed in the **Pain Management Protocol**.
 - c. For severe pain (greater than 7 on the Wong Pain Scale), administer opioid as directed in the **Pain Management Protocol**. Administer IV/IO slowly, titrating to lowest dose needed to reduce pain score to less than 7.
 - d. Opioids should not be used for labor pain.

Contraindications:

1. Hypotension
2. Children $\leq$ 18 months old

Expected effects:

1. Decreased pain

Side effects:

1. Respiratory depression
2. Hypotension

Dosing: PAIN MANAGEMENT

Adults administer:

1. Morphine 0.1 mg/kg IV/IO/IM. Maximum single dose 5 mg.
 - a. IV/IO may repeat three times (total of four doses). Total dose may not exceed 20 mg. Interval between doses is minimally 10 minutes.
 - b. If IM administration may NOT repeat.

Pediatrics (patients > 18 months of age) administer:

1. Morphine according to MI MEDIC
2. When MI MEDIC is not available administer Morphine 0.1 mg/kg IV/IO/IM. Maximum single dose 5 mg.
 - a. IV/IO may repeat three times (total of four doses). Total dose may not exceed 20 mg. Interval between doses is minimally 10 minutes.

Michigan
MEDICATION SECTION
MORPHINE

Initial Date: 07/19/2023

Revised Date: 02/10/2026

Section: 9-35R

- b. If IM administration may NOT repeat.
- c. Do NOT administer to children < 18 months of age.

Used in the Following Protocol(s):

Pain Management (Section 7 Procedures)